

The Cosmic Resilience- Solar Elastosis

Anubha Bajaj*

Consultant Histopathology, A.B. Diagnostics, New Delhi, India

Introduction

Solar elastosis occurs due to chronic, persistent cutaneous damage arising on account of ultraviolet radiation affecting ageing cutaneous surfaces. Solar elastosis exhibits an accumulation of anomalous elastin or elastic tissue within cutaneous dermal layer or ocular conjunctiva. The exceptional solar elastosis engendered by actinic-induced cutaneous damage exhibits thickened, wrinkled, dry, furrowed, bumpy and yellowish cutaneous surfaces (Figure-1). Solar elastosis is also designated as actinic elastosis or elastosis senilis [1,2].




Figure 1: Solar elastosis exhibiting a wrinkled, corrugated, bumpy, dry, yellowish cutaneous patches (11).

Preliminary lesions of solar elastosis demonstrate dermal proliferation of elastic fibres. Gradually, papillary and reticular dermal collagen is replaced by thick, curled, basophilic fibres configuring tangled masses. Thus, solar elastosis may be contemplated as a biological indicator and a protective, compensatory reaction to proportionate, ultraviolet-induced cutaneous injury. Confirmation by cogent histological assessment is mandated.

Disease Characteristics

Solar elastosis emerges as a feature of photo-ageing and occurs in individuals with extended exposure to sun. The lesion initially appears beyond > 40 years. Elderly subjects depicting solar elastosis are associated with enhanced possible emergence of cutaneous malignant melanoma [3,4].

Although devoid of racial predilection, incriminated Caucasian population predominantly exhibits a yellow cutaneous hue. Immuno-compromised individuals or Caucasians with minimally tinted irises and blonde hair subjected to excessive exposure to sun or pursuing an outdoor occupation depict a predilection for solar elastosis, rapidly emerging freckles or sunburn [3,4].

Accumulation and denaturation of elastic tissue ensues on account of cumulative, prolonged and excessive actinic exposure which induces photo-ageing. (Figure 2) Extended exposure to sunlight engenders solar elastosis wherein no cutaneous subtype is spared [3,4].

***Corresponding author:** Anubha Bajaj, Consultant Histopathology, A.B. Diagnostics, New Delhi, India, E-mail: anubha.bajaj1@outlook.com

Received Date: June 01, 2022

Accepted Date: June 06, 2022

Published Date: June 11, 2022

Citation: Anubha B (2022) The Cosmic Resilience- Solar Elastosis, J Clinic Exper Cosme Derma 5: 022.

Copyright: © 2022 Anubha B. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 2: Solar elastosis exemplifying degeneration of dermal elastic layer with a specific Grenz zone and epidermal acanthosis(12).

Solar elastosis engendered by smoking is indicative of premature cutaneous ageing. Tobacco consumption alters collagen production with elevated configuration of tropo-elastin and matrix metalloproteinases (MMPs) [3,4].

In fact, smoking enhances production of tropo-elastin fibres along with degradative enzymes constituted by Matrix Metallo Proteinases (MMPs). Consequently, matrix proteins are degraded with dermal accumulation of anomalous elastic substance. Alteration within elastic fibres may extend towards deep-seated reticular dermis [3,4].

Alternatively, singular actinic-induced damage in the absence of tobacco consumption is associated with alterations of elastic fibres predominantly restricted to superficial papillary dermis [3,4].

Of obscure aetiology, configuration and accumulation of elastotic substance within the dermis is posited to arise due to actinic stimulation of fibroblasts, a process which stimulates synthesis of the substance. Besides, elastotic substance emerges as a degradation product of collagen and elastin [3,4].

Dermal elastotic substance is posited to arise from degradation of collagen and elastic fibres, fibroblasts activated due to actinic induced injury with syntheses of fresh elastic tissue and an anomalous, synthetic, degradative tissue process [3,4].

Clinical Elucidation

Solar elastosis is associated with minimal clinical symptoms and may exemplify a gradually enlarging, reddish, bumpy cutaneous patch with pruritus or burning sensation. Solar elastosis initially incriminates sun-exposed cutaneous zones as the face, lips, ears, neck, forearms or hands. (Figure 3) Upon examination, a reddish or yellow papule of variable magnitude is observed [4,5].




Figure 3 : Solar elastosis enunciating elastotic degeneration separated by a definitive grenz zone and acanthosis of superimposed epidermis (12).

Accumulation of deranged elastic tissue ensues within actinic-sensitive upper and mid-dermis. Photo-ageing is associated with cutaneous damage confined to superficial papillary dermis [4,5].

Tobacco smoke induces denaturation of deep-seated reticular dermis. Tropo-elastin and matrix metalloproteinases (MMPs) denature the matrix with consequent, disorderly configuration of elastin fibres within cutaneous layers [4,5].

Solar elastotic syndrome manifests as thickened, dry, coarse, wrinkled cutaneous soft tissue associated with decimated skin tone [4,5].

The commonest clinical variant appears as thickened, intensely fissured cutaneous surface discerned upon dorsum of neck subjected to chronic actinic-induced damage which is designated as cutis rhomboidalis nuchae and configures a segment of cutaneous photo-ageing. Cutis rhomboidalis nuchae exhibits thickened, yellow, leathery cutaneous tissue confined to posterolateral neck surface [4,5].

Papular elastosis as a terminology was initially adopted by Kwitken wherein asymptomatic, solitary or multiple, shiny, smooth, firm, variably hued papules of magnitude up to 10 millimetres are associated with severe elastosis within the papillary cutis [6] (Figure 4).

Elastotic nodules confined to ear characteristically display singular or multiple, bilateral, firm or hard, pale nodules situated upon anti-helix or helix of the pinna. Upon histology, irregular, coarse, elastotic substance and clumped, thick fibres are admixed with foci of prominent solar elastosis and are appropriately appreciated with elastic-Van Gieson stain. Clinical segregation is required from basal cell carcinoma, amyloidosis, gouty tophi and chondrodermatitis nodularis helioides.




Figure 4 : Solar elastosis depicting significant acanthosis, denaturation of dermal elastic layer, a normal layer of collagen and mild, chronic inflammatory infiltrate (13).

Collagenous and elastotic plaques of the hand appear as a gradually progressive disorder discerned in elderly males. Lesions appear as waxy, linear plaques and are situated upon the junction of dorsal and palmar cutaneous surface of the hand [5,7].

Upon histology, amorphous, basophilic nodules of elastotic tissue and thickened, fragmented, calcified, elastic fibres are admixed within the upper and mid-dermis. Thickened, haphazard collagen bundles appear perpendicular to the cutaneous surface [5,7].

Adult colloid milium is a solar elastotic dermatosis which resembles papular solar elastosis. Upon microscopy, homogenous, amorphous, mildly eosinophilic substance is admixed with cleft-like spaces confined to upper and mid-dermis [5,7].

Of obscure aetiology, dermal colloid deposition ensues due to actinic-induced degeneration of collagen and elastic fibres. Besides, the elastotic substance may be engendered due to actinically injured fibroblasts [5,7].

Histological Elucidation

Solar elastosis can be appropriately discerned by cogent tissue sampling of incriminated zone. Decimated staining with eosin within upper dermal layers is associated with a basophilic hue and significant disorganization of dermal elastin fibres [7,8].

Solar elastosis is indicative of pertinent histopathological alterations with degeneration of dermal elastic tissue which ensues within photo-damaged cutaneous surfaces. Predominantly, solar elastosis exhibits basophilic degeneration of dermal elastotic fibres with a narrow band of normal, horizontally disseminated collagen or Grenz zone segregating the dermis from epidermis. Elastosis is proportionate to cumulative exposure of ultraviolet radiation to cutaneous surfaces (Figure 5).

Verhoeff's stain can be employed to stain the elastotic fibres black [7,8].

Upon microscopy, decimated eosin staining is observed within superficial dermis along with accumulation of irregular, dense elastic fibres. The elastic fibres are degraded to configure tangled tropoelastin and fibrillin articulations [7,8].

Foci of solar elastosis may be admixed with patchy, chronic, non-specific lymphocytic and plasma cell inflammatory infiltrate. Epidermal acanthosis may accompany the lesions [7,8].




Figure 5 : Solar elastosis displaying denatured dermal elastic layer, horizontal dissemination of collagen fibres, numerous congested vascular articulations and an acanthotic epidermal layer (14).

Additionally, Favre-Rachouchot syndrome is associated with elevated elastosis of upper and mid-dermis configuring thickened, tortuous elastic tissue fibres [7,8].

Differential Diagnosis

Papules of solar elastosis require segregation from lesions such as basal cell carcinoma, sebaceous hyperplasia, molluscum contagiosum or actinic keratosis [3,4].

Solar elastosis upon the hands requires demarcation from keratolastosis marginalis which is an acquired marginal keratoderma incriminating lateral aspect of index finger or thumb [3,4].

Therapeutic Options

Solar elastosis can be minimized with circumvention of sun exposure which is an efficacious prevention strategy. Adoption of appropriate sun protection with broad spectrum sunscreens and suitable clothing is recommended. Also, discontinuation of smoking and preventing exposure to passive smoking appears advantageous. Antecedent disease prevention and therapy is efficacious and recommended [9,10] (Figure 6).




Figure 6 : Solar elastosis demonstrating degeneration of dermal elastic layer, distinct grenz zone and acanthosis of superimposed epidermis with hyperkeratosis (15).

Treatment options of variable efficacy applicable for alleviating modifications of photo-aged cutaneous surfaces are comprised of dermabrasion, topical retinoic acid, topical oestrogens, resurfacing with carbon dioxide laser, dermal injections of hyaluronic acid, imiquimod or tacrolimus ointment [9,10].

Cosmetic outcomes of elastotic cutaneous surfaces can be meliorated with employment of laser ablation, topical imiquimod, injectable botulinum toxin which decimates nervous activity with cutaneous relaxation or dermal fillers which permeate cutaneous contours and decrease depth of cutaneous furrows [9,10].

Additionally, ablative and nonablative laser therapy, dermal fillers and injectable neurotoxins as botulinum toxin can be adopted in order to enhance aesthetic outcomes of solar elastosis therapy [9,10].

It is posited that imiquimod therapy may ameliorate morphological countenance of solar elastosis in accompaniment with elastotic dermal modifications [9,10].

Certain lesions may progress into cutaneous squamous cell carcinoma. Thus, annual monitoring of solar elastosis is mandated [9,10].

Organ transplant recipients on immunosuppressive therapy exhibit possibility of actinic damage and emergence of diverse cutaneous malignancies. Besides, deceleration of cutaneous carcinogenesis may ensue with discontinuation of immune-suppressive therapy [9,10].

References

1. Kligman AM (1969) Early destructive effect of sunlight on human skin. *JAMA* 210: 2377-2380.
2. Izumi AK, Marples RR (1973) Senile (solar) comedones. *J Invest Dermatol* 61: 46-50.
3. Heng JK, Aw DC (2014) Solar elastosis in its papular form: Uncommon, mistakable. *Case Rep Dermatol* 6: 124-128.
4. Yeh C, Schwartz RA (2021) Favre-Racouchot disease: Protective effect of solar elastosis. *Arch Dermatol Res* 314: 217-222.
5. Lewis KG, Bercovitch L (2004) Acquired disorders of elastic tissue. 1. Increased elastic tissue and solar elastotic syndromes. *J Am Acad Dermatol* 51: 1-21.
6. Kwittken J (2000) Papular elastosis. *Cutis* 66: 81-83.
7. Calderone DC, Fenske NA (1995) The clinical spectrum of actinic elastosis. *J Am Acad Dermatol* 32: 1016-1024.
8. Miri K, Jeong HK (2016) Molecular Mechanisms of Skin Aging and Rejuvenation. *Molecular Mechanisms of the Aging Process and Rejuvenation*.
9. Smith K, Hamza S (2007) Does imiquimod histologically rejuvenate ultraviolet radiation-damaged skin?. *Dermatol Surg* 33: 1419-1428.
10. Widge AD, Napkoski K (2021) New approaches to skin photodamage histology-Differentiating 'good' versus 'bad' Elastin. *J Cosmet Dermatol* 20: 526-531.



Henry Journal of Acupuncture & Traditional Medicine

Henry Journal of Anesthesia & Perioperative Management

Henry Journal of Aquaculture and Technical Development

Henry Journal of Cardiology & Cardiovascular Medicine

Henry Journal of Case Reports & Imaging

Henry Journal of Cell & Molecular Biology

Henry Journal of Tissue Biology & Cytology

Henry Journal of Clinical, Experimental and Cosmetic Dermatology

Henry Journal of Diabetes & Metabolic Syndrome

Henry Journal of Emergency Medicine, Trauma & Surgical Care

Henry Journal of Haematology & Hemotherapy

Henry Journal of Immunology & Immunotherapy

Henry Journal of Nanoscience, Nanomedicine & Nanobiology

Henry Journal of Nutrition & Food Science

Henry Journal of Obesity & Body Weight

Henry Journal of Cellular & Molecular Oncology

Henry Journal of Ophthalmology & Optometry

Henry Journal of Perinatology & Pediatrics

Submit Your Manuscript: <https://www.henrypublishinggroups.com/submit-manuscript/>

Henry Publishing Groups, 41341 Red Birch Dr, Aldie, VA, 20105, USA

Tel: +1 571-275-4480; E-mail: contact@henrypublishinggroups.org

<https://www.henrypublishinggroups.com/>
